

# MoorMan's® Show Tec® Finisher Paylean® DBZ 11256PLL

► **MEDICATED** ◀

## A Complete Feed for Finishing Pigs Being Fed for Exhibition

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine, weighing not less than 150 lbs, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lb of gain prior to slaughter.

### ACTIVE DRUG INGREDIENTS

Ractopamine hydrochloride

9.0 gm/lb

An insect growth regulator, which prevents the development of manure-breeding flies in the manure of treated swine (including nursery, grow-finish, and breeding stock)

**ACTIVE INGREDIENT:**

Diflubenzuron (CAS # 35367-38-5)

0.001%

**OTHER INGREDIENTS:**

99.999%

Contains 4.635 mg of diflubenzuron per lb.

Total 100.000%

GUARANTEED ANALYSIS	INGREDIENTS
Crude Protein, Min. .... 18.0%	Ground Corn, Dehulled Soybean Meal, Soybean Hulls, Sodium Bentonite, Lignin Sulfonate, Monocalcium Phosphate, Dicalcium Phosphate, Fish Meal, Meal and Bone Meal, Calcium Carbonate, Salt, Blood Meal, Calcium Propionate (A Preservative), Dried Whey, Copper Sulfate, Natural and Artificial Flavors, Extracted Citric Acid Presscake, Maltodextrin, Yeast Culture (Saccharomyces cerevisiae), Diatomaceous Earth, L-Lysine, Oxidized Chloride, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Zinc Oxide, Ferrous Sulfate, Vitamin E Supplement, Biotin, Manganese Sulfate, Zinc Sulfate, Nicotin Supplement, Calcium Panthothenate, Magnesium Oxide, Vitamin A Supplement, Menadione Dimethylpyrimidinol Bisulfite, Riboflavin Supplement, Vitamin B <sub>1</sub> Supplement, Vitamin B <sub>6</sub> Supplement, Calcium Iodide, Sodium Selenite
Lysine, Min. .... 1.0%	
Crude Fat, Min. .... 2.5%	
Crude Fiber, Max. .... 4.0%	
Calcium (Ca), Min. .... 0.7%	
Calcium (Ca), Max. .... 1.2%	
Phosphorus (P), Min. .... 0.6%	
Salt (NaCl), Min. .... 0.2%	
Salt (NaCl), Max. .... 1.7%	
Selenium (Se), Min. .... 0.3 ppm	
Zinc (Zn), Min. .... 155 ppm	
Biotin, Min. .... 0.14 mg/lb	

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

### USE AND MIXING DIRECTIONS

**STORAGE AND DISPOSAL:** Do not contaminate water or food by storage or disposal. **STORAGE:** Store in a cool, dry place. Do not contaminate with pesticides or fertilizer. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### FEEDING DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product only prevents the development of house flies, stable flies, face flies, horn flies and manure breeding flies in the manure of treated swine, but is not effective against existing adult flies. Begin use early in the spring, 30 days before flies begin to appear, and continue feeding throughout the summer and into the fall until cold weather restricts fly activity. In some cases, supplemental fly control measure may be needed in and around pastures, barns, and feedlots to control adult house and stable flies which can breed in other decaying, organic matter or sludge in the area, or that migrate in from other locations. MoorMan's Show Tec Finisher Paylean is a complete feed specifically formulated for pigs fed for exhibition. Feed continuously as the sole ration to finishing swine weighing not less than 150 lb for the last 45 to 90 lb (group average) of weight gain prior to slaughter. Feed at the rate of 5.0 lbs per head per day to animals weighing 250 lbs to provide 0.2 mg diflubenzuron per kg of body weight per day. The higher inclusion rate for diflubenzuron (0.2 to 0.3 mg/kg allowed) is to be used for lighter weight pigs. No increased benefit has been shown when Ractopamine concentrations in the diet are greater than 4.5 g/lb. Pigs must be fed in an ad libitum (full feed) feeding program. Provide clean, fresh water at all times.

**CAUTION:** Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for animals intended for breeding. Do not use in any finished feed containing in excess of 2% bentonite.

**WARNING:** The active ingredient in Paylean, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention.

\*Paylean is a registered trademark of Elanco, Indianapolis, IN

05124

ADM Animal Nutrition, a division of  
Archer Daniels Midland Company,  
Quincy, IL 62305-3115



7 99393 00277

6

219601NE

00926PMI

Net Weight 50 lb (22.67 kg) Or As Invoiced

11256PLLE4