

MoorMan's® ShowTec® Finisher Paylean® DBZ 11256PLL

MEDICATED

A Complete Feed for Finishing Pigs Bleag Fed for Eahbition

For increased rate of weight gain, improved feed efficiency and increased carcass, conness, in finishing swine, weighing and less than 150 lbs, fed a complete ration containing at least 15% crude protein for the just 45 to 60 IU of gain prior to slaughter.

ACTIVE DRUG INGREDIENTS

Ractopamine hydrochloride: 30.00%

An insect growth regulator, which prevents the development of muscoid breeding flies in the finisher of finished swine (including mature, grower, and breeding stock).

ACTIVE INGREDIENT:

Diflubenzuron (CAS # 9597-34-5)

0.00%

OTHER INGREDIENTS:

Total 0.00%

GUARANTEED ANALYSIS (continued)	INGREDIENTS
Lysine, Protein, Min, ANALYSES	
Lysine, Min. 0.0%	Ground Corn, Distilled Dried Beans, Soybean Hulls, Codon,
Lysine, Min. 1.5%	Conbottle, Lignin, Sulphuric, Monocyclic Phosphate, Sodium
Crude Fat, Min. 2.5%	Dried Yeast, Dried Molasses, Molasses, Molasses, Molasses,
Crude Fiber, Max. 3.0%	Salt, Dried Molasses, Calcium Phosphate (A Phosphate), Dried Yeast,
Calcium (Ca), Min. 0.7%	Crystalline Diphosphate, Sodium and Artificial Flavors, Edible Oil, Acid
Calcium (Ca), Max. 1.2%	Preservatives, Monocyclic Phosphate, Yeast, Calcium, (Calcium Phosphate)
Phosphorus (P), Max. 0.6%	Geraniol, Geraniol, Geraniol, Geraniol, Geraniol, Geraniol, Geraniol,
Serine (Ser), Min. 0.2%	Anaerobic Acid, Anaerobic, Anaerobic, Anaerobic, Anaerobic, Anaerobic,
Salt (NaCl), Max. 1.7%	Ferrous, Sulfate, Vitamin E, Supplementation, Fecund, Mangroves, Sulfate,
Selenium (Se), Min. 0.3 ppm	Zinc Sulfate, Niacin, Supplementation, Calcium Phosphate,
Zinc (Zn), Min. 0.3 mg/lb	Magnesium Oxide, Vitamin A Supplementation, Manganese,
Moisture, Min. 0.14 mg/lb	Dihydroxyacetone Phosphate, Phosphate, Phosphate, Phosphate, Phosphate,
	Supplementation, Vitamin B, Supplementation, Calcium Iodide, Sodium
	Sulfate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

USE AND MIXING DIRECTIONS

STORAGE AND DISPOSAL: Do not contaminate water or food by storage or disposal. **STORAGE:** Store in a cool, dry place. Do not contaminate with pesticides or fertilizers. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. If not disposed of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FEEDING DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product only prevents the development of house flies, stable flies, face flies, horn flies and midges breeding flies in the manure of healthy swine, fed in a complete ration. It does not affect the growth rate of the pig. In the spring, 30 days before flies begin to appear, and continue feeding throughout the summer and into the fall until cold weather restricts fly activity. In some cases, supplemented by control measures may be needed in and around pastures, barns, and feedlots to control adult house flies and stable flies which can breed in other decaying organic matter or sludge in the area, or feed material from other locations. **Showtec Finisher Paylean** is a complete feed specifically formulated for pigs, fed for exhibition. Feed continuously as the sole ration to finishing swine weighing and less than 150 lbs. The feed is to be fed (average) of weight gain prior to slaughter. Feed at the rate of 5.0 lbs. per head per day to animals weighing 250 lbs. to provide 12.2 mg diflubenzuron per kg of body weight per day. The highest feed rate may be diflubenzuron (0.28 to 0.3 mg/kg allowed) is to be fed for 15 lbs. weight pigs. No increased benefits have been shown when Ractopamine concentrations in the diet are greater than 15.3 g/lb. Pigs must be fed in a ration and a complete feeding program. Provide clean, fresh water at all times. **CAUTION:** Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Do not use in any feedlot or breeding herd containing in excess of 2% benzocaine. **WARNING:** The active ingredient in Paylean, Ractopamine, hydrochloride, is a beta-adrenergic agonist. It causes, with cardiovascular effects, skeletal tremors, spread cardiac to avoid exposure. Not to be used in humans. Keep out of the reach of children. Operators should wear thoroughly with soap and water, after handling. If contacted by or inhaled occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention.

*Paylean is a registered trademark of Eanco, Indianapolis, IN

16-14

ADM Animal Nutrition, a division of
Archer Daniels Midland Company
Quincy, IL 62406-0001

Net Weight 50 lb (22.67 kg) CH/BE Marked



219601NE

201926PM

1125i6PLLE4